

ON THE ANALYSIS OF TUBERCULOSIS STUDIES WITH INTERMITTENT MISSING SPUTUM DATA

BY DANIEL SCHARFSTEIN^{*}, ANDREA ROTNITZKY[†], MARIA ABRAHAM[‡]
AIDAN McDERMOTT^{*} RICHARD CHAISSON^{*} AND LAWRENCE GEITER[§]

Johns Hopkins University^{}, Universidad Torcuato Di Tella[†], Statistics
Collaborative[‡], Otsuka Novel Products[§]*

In randomized studies evaluating treatments for tuberculosis (TB), individuals are scheduled to be routinely evaluated for the presence of TB using sputum cultures. One important endpoint in such studies is the time of culture conversion, the first visit at which a patient's sputum culture is negative and remains negative. This paper addresses how to draw inference about treatment effects when sputum cultures are intermittently missing on some patients. We discuss inference under a novel benchmark assumption and under a class of assumptions indexed by a treatment-specific sensitivity parameter that quantify departures from the benchmark assumption. We motivate and illustrate our approach using data from a randomized trial comparing the effectiveness of two treatments for adult TB patients in Brazil.

1. Introduction. In the design of randomized studies evaluating competing treatments for patients with tuberculosis (TB), it is common to culture sputum for the presence of TB at regularly scheduled clinic visits over a specified time horizon. A primary goal in such studies is to estimate the treatment-specific distribution of the visit of first culture conversion [European Medicines Agency, Committee for Medicinal Products for Human Use (2010)], which we refer to as the time of culture conversion. Culture conversion is said to have occurred for a patient at a given visit, if the sputum cultures for that visit and all subsequent visits are negative. A key complication in the analysis is that culture results are missing at some visits, because the culture was contaminated, the patient could not produce sputum, or the patient did not show up. Culture conversion status at a given visit is unknown when from that visit onward at least one culture result is missing *and* all recorded culture results are negative. For a given patient, the set of visits with unknown culture conversion status is empty, or it consists of either a single visit or a set of consecutive visits. If the set is not empty, the time of culture conversion will be known to lie in an interval; however,

Keywords and phrases: Culture Conversion, Curse of Dimensionality, Exponential Tilt-
ing, Reverse-Time Hazard, Sensitivity Analysis

the time may not be interval censored in the classical sense. This is because certain data configurations may imply that the time of culture conversion cannot occur at certain visit times within the interval. To distinguish this data structure from classic interval censoring, we refer to the set of feasible times that are compatible with an individual’s data as the coarsening set. The coarsening set can include a single point or a set of points.

The treatment-specific distribution of time of culture conversion is not identified without untestable assumptions about the distribution of culture conversion status within the coarsening sets. There are countless ways of imposing such assumptions. The “worst-case” and “best-case” assumptions, leading to bounds on the treatment-specific distribution of time of culture conversion, are that the missing culture associated with the latest visit time at which culture conversion status is unknown is positive and the missing cultures associated with all visit times at which culture conversion status is unknown are negative, respectively. In considering alternative assumptions it is natural to condition on as much of the relevant data as possible. In addition to conditioning on observed culture results, it is natural to condition on auxiliary factors that are associated with the unknown results inside the coarsening set.

Most TB studies collect a key auxiliary time-varying variable. Sputum specimens are also evaluated by smear, and the results of the smear may be available when a culture is contaminated. Sputum smear is a less reliable assessment of clinical tuberculosis than sputum culture. It relies on the visualization of the bacteria through a microscope after staining of sputum with dyes that allows the microscopist to see so-called acid-fast TB bacteria. Sensitivity of the sputum smear is about 0.5, but it can vary by staining method used and clinical population. In contrast, sputum culture involves breaking down the sputum (which is very viscous), decontaminating the specimen to kill bacteria other than the mycobacteria, and inoculating it on culture media where the bacilli can grow. Sensitivity of the sputum culture is about 0.8-0.85 ([American Thoracic Society, 2000](#)). Roughly 65% of those with a positive culture are expected to have a positive smear, and nearly 100% of those with a positive smear are expected to have a positive culture ([American Thoracic Society, 2000](#)).

Another important auxiliary variable is baseline cavitation status. Patients with pulmonary tuberculosis who have cavities seen on a chest radiograph (cavitary tuberculosis) are more likely to have positive sputum smears, as they harbor larger numbers of tubercle bacilli than patients without cavities. It generally takes longer for patients with cavitary tuberculosis to convert their smears and cultures to negative during treatment, as they

have a larger bacterial load and the therapy must kill more organisms.

This article is motivated by data from a randomized TB study previously analyzed by Conde *et al.* (2009).¹ This phase II, double-blind, randomized trial compared moxifloxacin vs. ethambutol in adults with smear-positive tuberculosis at baseline in a hospital in Rio de Janeiro, Brazil. All 170 patients randomized (85 to each treatment arm) into the study were treated with a background regimen of isoniazid, rifampicin and pyrazinamide. Patients with negative, contaminated or drug-resistant *Mycobacterium tuberculosis* at baseline were excluded from the analysis, resulting in an analysis sample of 74 and 72 patients in the moxifloxacin and ethambutol groups, respectively. Treatment was scheduled to be given five days per week and was to be directly observed by study personnel. Smear and culture testing of sputum specimens (spontaneous or induced) was scheduled on specimens collected at baseline and every week for 8 weeks. In this study, 55.4% and 62.5% of patients in the moxifloxacin and ethambutol arms, respectively, had complete culture data through week 8. Time of culture conversion could be determined for 64.9% and 72.2% of patients in these arms; the remaining patients had their time of culture conversion coarsened.

In this article, we develop a method that estimates the treatment-specific distribution of time of culture conversion under a class of assumptions on the distribution of the coarsened time of culture conversion that conditions on *all of the relevant available data*, including sputum cultures, sputum smears and baseline data. Each assumption in the class is indexed by a treatment-specific sensitivity-analysis parameter which quantifies the magnitude of discrepancy from a specific benchmark assumption. In Section 2, we provide a preview of our proposed method. In Section 3, we introduce the data structure and notation. Section 4 discusses our modeling assumptions and the approach to inference. Section 5 presents an analysis of data from the Conde *et al.* study. The article concludes with a short discussion.

2. Preview. Our approach starts by imposing non-testable assumptions that identify the conditional distribution of time to culture conversion given the data, for every data configuration for which time of culture conversion is unknown. The crucial methodological challenge then is to make sensible identifying assumptions. Because time of culture conversion is determined by the results of sputum cultures, these assumptions ultimately identify the visit-specific probabilities of the last positive sputum culture

¹The dataset provided to us differs slightly from that of Conde *et al.* (2009). There are small differences in the number of observed cultures and the number of observed negative cultures at each week. All analyses reported in this paper are based on the data provided to us.

TABLE 1

Example of patient data. - denotes negative culture, + denotes positive culture, ? denotes unknown, green ? denotes a culture result that is irrelevant for determining culture conversion, red ? denotes a culture result that is relevant for determining culture conversion.

Line		Visit								Coarsening Set
		1	2	3	4	5	6	7	8	
1	Culture Results	?	+	?	-	?	-	-	-	
2	Conversion?	N	N	?	?	?	Y	Y	Y	{3, 4, 6}
3	Culture Results	?	+	?	-	+	-	-	-	
4	Conversion?	N	N	N	N	N	Y	Y	Y	{6}
5	Culture Results	?	+	?	-	-	-	-	-	
6	Conversion?	N	N	?	Y	Y	Y	Y	Y	{3, 4}
7	Culture Results	?	+	+	-	-	-	-	-	
8	Conversion?	N	N	N	Y	Y	Y	Y	Y	{4}
9	Culture Results	?	+	-	-	-	-	-	-	
10	Conversion?	N	N	Y	Y	Y	Y	Y	Y	{3}

given the data. Lack of culture conversion at a given visit can be determined without full knowledge of all subsequent results if the culture at that visit is positive or at least one subsequent culture is observed to be positive. It turns out that with “proper book-keeping,” we can achieve identifiability by imposing assumptions that suffice to identify the distribution of time of culture conversion but do not fully identify the joint distribution of results across visits. These conditions identify the reverse-time conditional hazards of time of culture conversion. This section illustrates these issues by means of an example.

The sputum culture data collected on one patient in the study illustrate the coarsening structure of culture conversion status and the time of culture conversion. The sputum culture data for this patient, we call Mary, are displayed in the first line of Table 1; the associated culture conversion statuses are displayed in line 2. Missing values are indicated by ?. Mary has negative cultures at visits 4, 6, 7 and 8 and a positive culture at visit 2. She is a culture converter at visits 6, 7 and 8 (labeled ‘Y’ in the second line), known not be a culture converter at visits 1 and 2 (labeled ‘N’ in the second line) and has unknown culture conversion status at visits 3, 4 and 5 (labeled ‘?’ in the second line). First, even though her culture status at visit 1 (labeled with a green ‘?’ in the first line) is missing, it is irrelevant for determining whether she is a culture converter at that visit, because she has a positive culture at visit 2 and therefore cannot be a culture converter at visit 1. Second, missingness of cultures at visits 3 and 5 (labeled with a red ‘?’ in the first line) affects our ability to determine her culture conversion status at visits

3, 4 and 5 because all cultures after visit 5 are negative and the culture at visit 4 is also negative. So even though Mary has a negative culture at visit 4, her culture conversion status is not known at that visit. Third, the visits at which culture conversion status is unknown are consecutive: the earliest and latest visits are 3 and 5, respectively. Finally, the coarsening set for time of culture conversion comprises visits 3, 4 and 6. This follows because if the culture at visit 5 were positive, then the time of culture conversion would be visit 6, and if the culture at visit 5 were negative, the time of culture conversion would be either visit 3 or visit 4, depending on the result at visit 3.

To illustrate our approach for identifying the conditional distribution of time of culture conversion given the data, consider subset A , the subset of patients with the same observed data as Mary. Specifically, we must model the probability that the cultures at visits 3 and 5 are both negative (in which case the time of culture conversion is visit 3), the probability that the culture at visit 3 is positive and the culture at visit 5 is negative (in which case the time of culture conversion is visit 4) and the probability that the culture at visit 5 is positive (in which case the time of culture conversion is visit 6). In modeling these probabilities, it is natural to consider two chronological factorizations. In forward time, we would need to model the probability of a negative culture at visit 3, the conditional probability of a negative culture at visit 5 given negative culture at visit 3, and the conditional probability of a negative culture at visit 5 given a positive culture at visit 3. In reverse time, we would need to model (i) the probability of a positive culture at visit 5 and (ii) the conditional probability of a positive culture at visit 3 given a negative culture at visit 5. We utilize this latter factorization as it requires fewer modeling assumptions.

We first turn to the task of imposing assumptions that identify (i). Our identifying assumption specifies that (i) is the same as the probability of a positive culture at visit 5 among patients with the same data as in subset A with the exception that they have an observed culture at visit 5. The observed culture results for these patients are depicted in lines 3 and 5. These patients have observed culture conversion status as depicted in lines 4 and 6. The probability (i) is then assumed to be equal to the ratio of the proportion of patients with observed results as in line 3 to the sum of the proportions of patients with observed results as in lines 3 and 5.

Next, we turn to the task of imposing assumptions that identify (ii). Our identifying assumption specifies that (ii) is the same as the probability of a positive result at visit 3 among patients with the same pattern of observed cultures as that in line 5, with the exception that they have an observed

culture at visit 3. The results of these patients are depicted in lines 7 and 9 with associated culture conversion status in lines 8 and 10. The probability (ii) is then assumed to be equal to the ratio of the proportion of patients with observed cultures as in line 7 to the sum of the proportions of patients with observed cultures as lines 7 and 9.

In datasets of typical size, we will not be able to obtain estimates of the proportion of patients who have specific patterns of observed data due to the curse of dimensionality. As a result, our inference will require dimension reduction assumptions. We will use fully parametric models for the treatment-specific distributions of the data. These models are described in Section 3.4.

In Section 3.3, we evaluate the sensitivity of our results to our identifying assumptions by conducting inference under a class of exponential tilt deviations from the assumed conditional probabilities for the unobserved culture conversion status.

3. Formalization of the problem. Since we focus on inference about the time of culture conversion, separately for each treatment arm, we consider, until Section 4, only data from one arm and suppress notational dependence on treatment assignment.

3.1. Data Structure and Notation. Let X denote baseline cavitation status (1 for cavitation, 0 otherwise), C_k denote the indicator that the culture is negative at visit k (1 for negative, 0 for positive) and S_k denote the indicator that the smear is negative at visit k (1 for negative, 0 for positive). Let M_k^c and M_k^s be the indicators that C_k and S_k are missing, respectively (1 for missing, 0 for observed). The data recorded on an individual at visit k are realizations of the random vector $O_k = (M_k^c, C_k^{obs}, M_k^s, S_k^{obs})$ where $C_k^{obs} = C_k$ if C_k is observed and C_k^{obs} is empty otherwise and S_k^{obs} is defined likewise.

Let K denote the number of scheduled post-baseline visits. For any collection of random vectors $\{W_k\}_{1 \leq k \leq K}$, we use the notation $\overline{W}_k = (W_1, \dots, W_k)$. With this notation, the data recorded on an individual throughout the entire study are realizations of the random vector $\mathbf{O} = (X, \overline{O}_K)$.

Define time of culture conversion T to be the earliest visit such that sputum cultures are negative from that visit onward if such a visit exists and $T = K + 1$ otherwise. As illustrated in Section 2, when the data available on each patient is the vector \mathbf{O} defined earlier, T belongs to a set with a single visit time (in which case T is determined from \mathbf{O}) or a set with multiple visit times, not necessarily consecutive. We denote the coarsening set where

T is known to lie by \mathcal{T} . Given \mathbf{O} , T is determined (i.e., the set \mathcal{T} has one element), unless either

- (i) the culture at visit K is missing, i.e. $M_K^c = 1$, or
- (ii) there exists a visit $k < K$ with a missing culture, i.e. $M_k^c = 1$, such that all subsequent visits have sputum cultures that are either negative or missing, i.e. $M_j^c = 1$ or $C_j^{obs} = 1$ for $j > k$.

When either (i) or (ii) occurs, then \mathcal{T} will have multiple visit times. We denote the lowest visit number in the set by L , which is the earliest visit k where the sputum cultures are either missing or negative at and subsequent to visit k . We denote the largest visit number in the set by $R + 1$, where $R = K$ if the culture at visit K is missing (i.e. $M_K^c = 1$) and $R = k$ ($k < K$) if at visit k the sputum culture is missing and at all subsequent visits the sputum cultures are recorded and negative. The other times in the coarsening set include all visit numbers k such that $L < k < R + 1$ and $M_{k-1}^c = 1$.

Formally, our inferential goal is to estimate the distribution of time of culture conversion, i.e., $P[T = k]$ for $k = 1, \dots, K$, based on n i.i.d. realizations of the vector \mathbf{O} . We use the subscript i to denote data for the i th individual.

In the next section we formally describe the identifying assumptions on which our benchmark analysis relies. These assumptions were illustrated in Section 2. The assumptions are “identifying” in the sense that once they are imposed, we are able to express $P[T = k]$ for $k = 1, \dots, K$ as a function of the distribution of the observed data \mathbf{O} and consequently, we can hope, to estimate $P[T = k]$ consistently. Subsequently, we propose models for departures from the benchmark assumptions that form the basis of our proposed sensitivity analysis. Specifically, our sensitivity analysis consists of repeating estimation of $P[T = k]$ under various plausible departures from the benchmark assumptions.

Both our benchmark analysis and the models for our sensitivity analysis rely on assumptions that identify the conditional distribution of T given the observed data \mathbf{O} . The marginal distribution of T is then obtained as the mixture of the, now identified, conditional distribution of T given \mathbf{O} mixed over the distribution of the observed data \mathbf{O} .

3.2. Benchmark Identifying Assumptions. To help guide our choice of benchmark identifying assumptions, we first note that since $P[T = k|\mathbf{O}] = 0$ if $k \notin \mathcal{T}$ and $P[T = k|\mathbf{O}] = 1$ if $k \in \mathcal{T}$ and $|\mathcal{T}| = 1$, we only need assumptions that suffice to identify $P[T = k|\mathbf{O}]$ when $k \in \mathcal{T}$ and $|\mathcal{T}| > 1$. Thus, we proceed in reverse order through the set \mathcal{T} by postulating assumptions that identify, first $P[T = R + 1|\mathbf{O}]$, then in reverse sequential order $P[T = k|\mathbf{O}] \leq$

$k, \mathbf{O}]$ where $k \in \mathcal{T}$, $L < k < R + 1$. This iterative procedure results in assumptions that identify $P[T = k|\mathbf{O}]$ for all $k \in \mathcal{T}$. This follows because $P[R + 1|\mathbf{O}]$ is identified and for $k \in \mathcal{T}, k < R + 1$, $P[T = k|\mathbf{O}]$ equals

$$P[T \neq R + 1|\mathbf{O}] \left\{ \prod_{\substack{k < s < R+1 \\ k \in \mathcal{T}}} P[T \neq s|T \leq s, \mathbf{O}] \right\} P[T = k|T \leq k, \mathbf{O}]$$

We need to focus only on the case where $|\mathcal{T}| > 1$. To guide our choice of benchmark assumptions for identifying $P[T = r + 1|\mathbf{O} = \mathbf{o}]$, we first note that given $\mathbf{O} = \mathbf{o}$ the event $T = r + 1$ occurs if and only if the culture at visit r is positive, i.e. if $C_r = 0$. Our benchmark assumption equates the unidentified probability $P[T = r + 1|\mathbf{O} = \mathbf{o}]$ with the identified probability that $C_r = 0$ in the subset of patients for which $\mathbf{O} = \mathbf{o}^{(r)}$ where $\mathbf{o}^{(r)}$ agrees with \mathbf{O} in all its components except that the culture at visit r is observed. Now, we write the event $\mathbf{O} = \mathbf{o}$ as the event

(3.1)

$$M_r^c = 1, \overline{M}_{r-1}^c = \overline{m}_{r-1}^c, \overline{C}_{r-1}^{obs} = \overline{c}_{r-1}^{obs}, V = v, M_j^c = 0, C_j = 1 \text{ for } r+1 \leq j \leq K$$

we define the event $\mathbf{O} = \mathbf{o}^{(r)}$ as the event

$$M_r^c = 0, \overline{M}_{r-1}^c = \overline{m}_{r-1}^c, \overline{C}_{r-1}^{obs} = \overline{c}_{r-1}^{obs}, V = v, M_j^c = 0, C_j = 1 \text{ for } r+1 \leq j \leq K$$

and we postulate that

$$(3.2) \quad P[T = r + 1|\mathbf{O} = \mathbf{o}] = P[T = r + 1|\mathbf{O} = \mathbf{o}^{(r)})$$

Next, we consider assumptions that identify $P[T = k|T \leq k, \mathbf{O} = \mathbf{o}]$ for $k \in \mathcal{T}, l < k < r + 1$. Once again, given $(T \leq k, \mathbf{O} = \mathbf{o})$, the event $T = k$ occurs if and only if the culture result at visit $k - 1$ is positive, i.e. $C_{k-1} = 0$. Because $k \in \mathcal{T}, l < k < r + 1$, we know that C_{k-1} is missing, i.e., $M_{k-1}^c = 1$. Our benchmark assumption in this case equates the probability $P[T = k|T \leq k, \mathbf{O} = \mathbf{o}]$ with the identified probability that $T = k$ (which equates to the event $C_{k-1} = 0$) in the subset of patients for which $\mathbf{O} = \mathbf{o}^{(k-1)}$, where $\mathbf{o}^{(k-1)}$ differs from the subset of patients with $(T \leq k, \mathbf{O} = \mathbf{o})$ only in that a sputum culture is observed at visit $k - 1$ and the event $T \leq k$ is observed to occur (i.e., $M_j^c = 0$ and $C_j = 1$ for all $k \leq j \leq K$). Formally, with the event $\mathbf{O} = \mathbf{o}$ defined as in (3.1), we define the event $\mathbf{O} = \mathbf{o}^{(k-1)}$ as the event

$$M_{k-1}^c = 0, \overline{M}_{k-2}^c = \overline{m}_{k-2}^c, \overline{C}_{k-2}^{obs} = \overline{c}_{k-2}^{obs}, V = v, M_j^c = 1, C_j = 1 \text{ for } k \leq j \leq K$$

and assume

$$(3.3) \quad P[T = k|T \leq k, \mathbf{O} = \mathbf{o}] = P[T = k|\mathbf{O} = \mathbf{o}^{(k-1)}]$$

Patients in the subset defined by $(T \leq k, \mathbf{O} = \mathbf{o})$ and subjects in the subset $\mathbf{O} = \mathbf{o}^{(k-1)}$ have the same baseline factors and the same recorded history of the auxiliary smear sputums throughout the study, as well as the same recorded history of sputum cultures up to visit $k - 2$.

Finally, for a realization $\mathbf{O} = \mathbf{o}$ where $|\mathcal{T}| > 1$, $P[T = l|T \leq l, \mathbf{O} = \mathbf{o}] = 1$.

3.3. Sensitivity Analysis. The benchmark assumptions (3.2) and (3.3) are untestable. For realizations $\mathbf{O} = \mathbf{o}$ with $|\mathcal{T}| > 1$, the following exponential tilt model [Cox and Barndorff-Nielsen (1994)] expresses departures from our benchmark assumptions:

$$(3.4) \quad P[T = r + 1|\mathbf{O} = \mathbf{o}] = \frac{P[T = r + 1|\mathbf{O} = \mathbf{o}^{(r)}] \exp(\alpha)}{h_{r+1}(\mathbf{o}^{(r)}; \alpha)}$$

and for $k \in \mathcal{T}$, $l < k < r + 1$,

$$(3.5) \quad P[T = k|T \leq k, \mathbf{O} = \mathbf{o}] = \frac{P[T = k|\mathbf{O} = \mathbf{o}^{(k-1)}] \exp(\alpha)}{h_k(\mathbf{o}^{(k-1)}; \alpha)}$$

where α is fixed and given and $h_k(\mathbf{o}^{(k-1)}; \alpha)$ are normalizing constants equal to $E[\exp\{\alpha I(T = k)\}|\mathbf{O} = \mathbf{o}^{(k-1)}]$ for $k \in \mathcal{T}$, $l < k \leq r + 1$. Under this exponential tilt model,

$$\frac{\text{odds}(P[T = r + 1|\mathbf{O} = \mathbf{o}])}{\text{odds}(P[T = r + 1|\mathbf{O} = \mathbf{o}^{(r)}])} = \frac{\text{odds}(P[T = k|T \leq k, \mathbf{O} = \mathbf{o}])}{\text{odds}(P[T = k|\mathbf{O} = \mathbf{o}^{(k-1)}])} = \exp(\alpha)$$

Thus, the magnitude of α quantifies the departure from our benchmark assumptions. When $\alpha > 0$ (< 0), $P[T = r + 1|\mathbf{O} = \mathbf{o}]$ is greater (less) than $P[T = r + 1|\mathbf{O} = \mathbf{o}^{(r)}]$ and $P[T = k|T \leq k, \mathbf{O} = \mathbf{o}]$ is greater (less) than $P[T = k|\mathbf{O} = \mathbf{o}^{(k-1)}]$. As $\alpha \rightarrow \infty$ ($-\infty$), $P[T = r + 1|\mathbf{O} = \mathbf{o}]$ and $P[T = k|T \leq k, \mathbf{O} = \mathbf{o}]$ go to one (zero). When $\alpha \rightarrow \infty$ ($-\infty$), the “worst-case” and “best-case” bounds described in the introduction are attained. $\alpha = 0$ corresponds to the benchmark assumption. To facilitate sensitivity analysis, our class of models assumes that the departures from the benchmark assumption are not time-specific.

3.4. Modeling. For specified α , estimation of the distribution of time of culture conversion depends on our ability to estimate for each realization

$O = o$ with $|\mathcal{T}| > 1$, $P[T = k | \mathbf{O} = \mathbf{o}^{(k-1)}]$ for $k \in \mathcal{T}, l < k \leq r + 1$. However, in practice these probabilities cannot be estimated non-parametrically. Therefore, we use a parametric model for the law of the observed data \mathbf{O} given baseline covariates X . This model induces parametric models for $P[T = k | \mathbf{O} = \mathbf{o}^{(k-1)}], k \in \mathcal{T}, l < k \leq r + 1$, that ultimately enable estimation of $P[T = k | \mathbf{O} = \mathbf{o}]$ by borrowing information across strata $\mathbf{O} = \mathbf{o}^{(k-1)}$.

We first recall that $\mathbf{O} = (X, \bar{O}_K)$, where $O_k = (M_k^c, C_k^{obs}, M_k^s, S_k^{obs})$ are the data available at visit k . We model the law of \mathbf{O} by modeling the distribution of O_k given \bar{O}_{k-1} and X for all $k = 1, \dots, K$, where $\bar{O}_0 = \emptyset$. We use separate logistic regression models for

1. the probability of $M_k^c = 1$ given \bar{O}_{k-1} and X , i.e.

$$(3.6) \quad \text{logit}\{P[M_k^c = 1 | \bar{O}_{k-1}, X]\} = a(k, \bar{O}_{k-1}, X; \gamma^{(a)})$$

2. the probability that $C_k^{obs} = 1$ given $M_k^c = 0, \bar{O}_{k-1}$ and X , i.e.

$$(3.7) \quad \text{logit}\{P[C_k^{obs} = 1 | M_k^c = 0, \bar{O}_{k-1}, X]\} = b(k, \bar{O}_{k-1}, X; \gamma^{(b)})$$

3. the probability that $M_k^s = 1$ given $M_k^c, C_k^{obs}, \bar{O}_{k-1}$ and X , i.e.

$$(3.8) \quad \text{logit}\{P[M_k^s = 1 | M_k^c, C_k^{obs}, \bar{O}_{k-1}, X]\} = c(k, M_k^c, C_k^{obs}, \bar{O}_{k-1}, X; \gamma^{(c)})$$

4. the probability that $S_k^{obs} = 1$ given $M_k^s = 0, M_k^c, C_k^{obs}, \bar{O}_{k-1}$ and X , i.e.

$$(3.9) \quad \text{logit}\{P[S_k^{obs} = 1 | M_k^s = 0, M_k^c, C_k^{obs}, \bar{O}_{k-1}, X]\} = d(k, M_k^c, C_k^{obs}, \bar{O}_{k-1}, X; \gamma^{(d)})$$

where $a(\cdot), b(\cdot), c(\cdot)$ and $d(\cdot)$ are specified functions of their arguments and $\gamma^{(a)}, \gamma^{(b)}, \gamma^{(c)}$ and $\gamma^{(d)}$ are unknown parameter vectors.

For the distributions in (3.7) and (3.9) we need not consider the cases $M_k^c = 1$ and $M_k^s = 1$, respectively, because for such settings the conditional distributions are degenerate (C_k^{obs} is empty when $M_k^c = 1$, and S_k^{obs} is empty when $M_k^s = 1$).

3.5. Inference. Under models (3.6) - (3.9) we can express for all realizations $\mathbf{O} = \mathbf{o}$ with $|\mathcal{T}| > 1$, the conditional probability $P[T = k | \mathbf{O} = \mathbf{o}^{(k-1)}]$ for $k \in \mathcal{T}, l < k \leq r + 1$ as a given function of $\mathbf{o}^{(k-1)}$ and $\gamma = (\gamma^{(a)}, \gamma^{(b)}, \gamma^{(c)}, \gamma^{(d)})$ whose expressions, for the special case where the right-hand sides of (3.6) - (3.9) only depend on O_{k-1} , are given in the Appendix. We denote this expression as $P[T = k | \mathbf{O} = \mathbf{o}^{(k-1)}; \gamma]$. Consequently, if we additionally assume models (3.4) and (3.5), then we can express $P[T = r + 1 | \mathbf{O} = \mathbf{o}]$ and $P[T = k | T \leq k, \mathbf{O} = \mathbf{o}]$ ($k \in \mathcal{T}, l < k \leq r$) as

given functions, $P[T = r + 1 | \mathbf{O} = \mathbf{o}; \gamma; \alpha]$ and $P[T = k | T \leq k, \mathbf{O} = \mathbf{o}; \gamma; \alpha]$ of o , γ and α .

The first step is to estimate γ using maximum likelihood; denote this estimator by $\hat{\gamma}$. This can be done using standard logistic regression software. This step does not rely on specification of the sensitivity analysis parameter α .

For fixed α , we estimate $P[T = k]$ by $\hat{P}[T = k; \alpha] = \frac{1}{n} \sum_{i=1} \hat{P}[T_i = k; \alpha]$ where $\hat{P}[T_i = k; \alpha]$ has one of four expressions depending of k and \mathbf{O}_i . If $k \notin \mathcal{T}_i$ then $\hat{P}[T_i = k; \alpha] = 0$; if $|\mathcal{T}_i| = 1$ and $k \in \mathcal{T}_i$ then $\hat{P}[T_i = k; \alpha] = 1$; if $|\mathcal{T}_i| > 1$ and $k = R_i + 1$ then $\hat{P}[T_i = k; \alpha] = P[T = R_i + 1 | \mathbf{O} = \mathbf{O}_i; \hat{\gamma}; \alpha]$; if $|\mathcal{T}_i| > 1$ and $k \in \mathcal{T}_i, k < R_i + 1$ then $\hat{P}[T_i = k; \alpha]$ equals

$$P[T \neq R_i + 1 | \mathbf{O} = \mathbf{O}_i; \hat{\gamma}; \alpha] \left\{ \prod_{\substack{k < s < R_i + 1 \\ k \in \mathcal{T}_i}} P[T \neq s | T \leq s, \mathbf{O} = \mathbf{O}_i; \hat{\gamma}; \alpha] \right\} \times \\ P[T = k | T \leq k, \mathbf{O} = \mathbf{O}_i; \hat{\gamma}; \alpha]$$

To compare the treatment-specific distributions of time to culture conversion, one can estimate a common treatment effect over time. Towards this end, one can utilize the logistic model for discrete survival data proposed by Cox (1972). This model assumes that

$$\frac{h_z(k)}{1 - h_z(k)} = \tau_k \exp(\beta z) \quad k = 1, \dots, K, z = 0, 1$$

where z denotes treatment group, $h_z(k) = P_z[T = k | T \geq k]$, and $\tau_1, \dots, \tau_K \geq 0$. Here $\exp(\beta)$ is the ratio of the odds of first becoming a culture converter at visit k given culture conversion at or after visit k , comparing moxifloxacin to ethambutol.

To estimate the model parameters, one can use equally weighted minimum distance estimation [Newey and McFadden (1994)]. Specifically, for each choice of α_z ($z = 0, 1$), we minimize the following objective function:

$$\sum_{z=0}^1 \sum_{k=1}^8 \left\{ \frac{\hat{h}_z(k; \alpha_z)}{1 - \hat{h}_z(k; \alpha_z)} - \tau_k \exp(\beta z) \right\}^2$$

with respect $\tau_1, \dots, \tau_8 \geq 0$ and β , where $\hat{h}_z(k; \alpha_z) = \hat{P}_z[T = k | T \geq k; \alpha_z]$. For each choice of α_z ($z = 0, 1$), this method finds the “closest” fitting logistic model to the “data”: $\{\hat{h}_z(k; \alpha_z) : k = 1, \dots, 8, z = 0, 1\}$. Even if the model is incorrectly specified, it can still be used to provide a valid test of the null hypothesis of no treatment effect.

To estimate the standard error of our estimator, we propose the use of non-parametric bootstrap.

4. Data Analysis. Figure 1 displays the treatment-specific observed culture results, with rows denoting patients, columns denoting visit number, red indicating a positive culture, green indicating a negative culture and white indicating a missing culture. Figure 2 displays the treatment-specific coarsening sets for time of culture conversion, with red and green indicating the infeasible and feasible points, respectively.

The treatment groups were not balanced with respect to cavitation status at baseline; 81.1% and 56.9% have cavitation in the moxifloxacin and ethambutol arms, respectively. It is essential that our analysis adjust for this key confounder. For each treatment group, we estimate the distribution of time of culture conversion by a weighted average of cavitation-specific distribution of time of culture conversion. The weights are taken to be the marginal (i.e., not conditional on treatment arm) proportion of patients with and without cavitation at baseline, respectively.

In our data analysis, we considered parsimonious models for the right-hand sides of (3.6) - (3.9). Our choice of models was guided by substantive considerations discussed with our scientist collaborators and by data analytic model fitting techniques. Our final model assumed that the right-hand sides of (3.6) - (3.9) depended only on $O_{k-1} = (M_{k-1}^c, C_{k-1}^{obs}, M_{k-1}^s, S_{k-1}^{obs})$ (i.e., not on the other components of \bar{O}_{k-1}) and the final model for (3.8) further assumed that $c(k, M_k^c, C_k^{obs}, O_{k-1}, X; \gamma^{(c)})$ did not depend on C_k^{obs} and C_{k-1}^{obs} . The latter assumption was imposed because missingness of a sputum culture at visit k is highly predictive of missingness of smear sputums at visit k and $k - 1$. In the Appendix we show that when the function $c(k, M_k^c, C_k^{obs}, O_{k-1}, X; \gamma^{(c)})$ does not depend on C_k^{obs} and C_{k-1}^{obs} for all k , then $P[T = k | O = o^{(k-1)}]$ does not depend on $c(k, M_k^c, C_k^{obs}, O_{k-1}, X; \gamma^{(c)})$ for any k , thus alleviating the need to further specify the function $c(k, M_k^c, C_k^{obs}, O_{k-1}, X; \gamma^{(c)})$. In the remaining models, we “borrowed strength”

FIG 1. Treatment-specific observed culture results, with rows denoting patients, columns denoting visit number, red indicating a positive culture, green indicating a negative culture and white indicating a missing culture.





FIG 2. *Treatment-specific coarsening sets, with rows denoting patients, columns denoting visit number, red indicating infeasible points and green indicating feasible points.*



across treatment groups. Specifically, we assumed

$$\begin{aligned}
a_z(k, O_{k-1}, X; \gamma^{(a)}) &= \gamma_{0,0,k}^{(a)} + \gamma_{0,1,k}^{(a)} X + \gamma_1^{(a)} I(k > 1) M_{k-1}^c + \gamma_2^{(a)} I(k > 1) (1 - M_{k-1}^c) C_{k-1}^{obs} + \\
&\quad \gamma_3^{(a)} I(k > 1) M_{k-1}^s + \gamma_4^{(a)} I(k > 1) (1 - M_{k-1}^s) S_{k-1}^{obs} + \gamma_5^{(a)} z \\
b_z(k, O_{k-1}, X; \gamma^{(b)}) &= \gamma_{0,k}^{(b)} + \gamma_1^{(b)} I(k > 1) M_{k-1}^c + \gamma_2^{(b)} I(k > 1) (1 - M_{k-1}^c) C_{k-1}^{obs} + \\
&\quad \gamma_3^{(b)} I(k > 1) M_{k-1}^s + \gamma_4^{(b)} I(k > 1) (1 - M_{k-1}^s) S_{k-1}^{obs} + \gamma_5^{(b)} z + \gamma_6^{(b)} X \\
d_z(k, M_k^c, C_k^{obs}, O_{k-1}, X; \gamma^{(d)}) &= \gamma_{0,k}^{(d)} + \gamma_1^{(d)} M_k^c + \gamma_2^{(d)} (1 - M_k^c) C_k^{obs} + \gamma_3^{(d)} (1 - M_k^c) C_k^{obs} X + \\
&\quad \gamma_4^{(d)} I(k > 1) M_{k-1}^c + \gamma_5^{(d)} I(k > 1) (1 - M_{k-1}^c) C_{k-1}^{obs} + \gamma_6^{(d)} I(k > 1) M_{k-1}^s + \\
&\quad \gamma_7^{(d)} I(k > 1) (1 - M_{k-1}^s) S_{k-1}^{obs} + \gamma_8^{(d)} z + \gamma_9^{(d)} X
\end{aligned}$$

where the functions are subscripted by treatment z ($z = 0$ denotes ethambutol, $z = 1$ denoted moxifloxacin).

In Tables 2, 3 and 4, we present estimates of the exponentiated parameters from these models, along with 95% non-parametric bootstrap percentile confidence intervals (based on 1000 resamples within treatment groups). In Table 2, missingness of sputum culture (aOR= 4.45; 95% CI: 2.03-9.07) and missingness of smear (aOR= 3.51; 95% CI: 1.52-9.66) at a previous visit are significant predictors of missingness of sputum culture at the next visit. In Table 3, among patients with an observed sputum culture at visit k , missingness of sputum culture (aOR = 4.00; 95% CI: 2.00-9.45), missingness of a smear culture (aOR = 5.16; 95% CI: 1.67-22.47), a negative observed culture (aOR = 6.37; 95% CI: 4.08-10.39) and a negative observed smear (aOR = 3.93; 95% CI: 2.51-6.18) at visit $k - 1$, as well as assignment to the moxifloxacin arm (aOR = 2.06; 95% CI: 1.46-3.19), are significant predictors of a negative sputum culture at visit k . In Table 4, among patients with an observed smear at visit k , missingness of smear (aOR = 3.66; 95% CI: 1.45-14.53) and an observed negative smear (aOR = 6.99; 95% CI: 4.50-11.81) at visit $k - 1$, as well as an observed negative sputum culture at visit k (aOR = 10.73; 95% CI: 5.58-31.18), are significant predictors of a negative smear at visit k .

Under our benchmark assumption, the estimated hazard ratio is 3.41 (95% CI: [1.33,13.06]), indicating that patients treated with moxifloxacin have a statistically significant shorter time of culture conversion than those treated with ethambutol. Figure 3 displays a contour plot of the estimated odds

TABLE 2

(Exponentiated) parameter estimates and 95% confidence intervals from the model for missingness of culture results. See $a_z(k, O_{k-1}, X; \gamma^{(a)})$ for form of the model.

Intercept	Odds	95% CI
Wk 1 ($\exp(\gamma_{0,0,1}^{(a)})$)	0.14	[0.04,0.30]
Wk 2 ($\exp(\gamma_{0,0,2}^{(a)})$)	0.11	[0.03,0.24]
Wk 3 ($\exp(\gamma_{0,0,3}^{(a)})$)	0.03	[0.00,0.09]
Wk 4 ($\exp(\gamma_{0,0,4}^{(a)})$)	0.11	[0.03,0.25]
Wk 5 ($\exp(\gamma_{0,0,5}^{(a)})$)	0.07	[0.01,0.19]
Wk 6 ($\exp(\gamma_{0,0,6}^{(a)})$)	0.04	[0.00,0.12]
Wk 7 ($\exp(\gamma_{0,0,7}^{(a)})$)	0.05	[0.00,0.14]
Wk 8 ($\exp(\gamma_{0,0,8}^{(a)})$)	0.07	[0.01,0.18]
Predictor	Odds Ratio	95% CI
Wk 1*Cav ($\exp(\gamma_{0,1,1}^{(a)})$)	0.19	[0.00,0.90]
Wk 2*Cav ($\exp(\gamma_{0,1,2}^{(a)})$)	0.21	[0.00,0.87]
Wk 3*Cav ($\exp(\gamma_{0,1,3}^{(a)})$)	2.93	[0.81, ∞] [†]
Wk 4*Cav ($\exp(\gamma_{0,1,4}^{(a)})$)	0.38	[0.09,1.50]
Wk 5*Cav ($\exp(\gamma_{0,1,5}^{(a)})$)	1.63	[0.55,11.28]
Wk 6*Cav ($\exp(\gamma_{0,1,6}^{(a)})$)	1.39	[0.36, ∞] [†]
Wk 7*Cav ($\exp(\gamma_{0,1,7}^{(a)})$)	1.60	[0.47, ∞] [†]
Wk 8*Cav ($\exp(\gamma_{0,1,8}^{(a)})$)	1.63	[0.51,8.32]
$I(k > 1)M_{k-1}^c$ ($\exp(\gamma_1^{(a)})$)	4.45	[2.03,9.70]
$I(k > 1)(1 - M_{k-1}^c)C_{k-1}^{obs}$ ($\exp(\gamma_2^{(a)})$)	0.89	[0.50,1.68]
$I(k > 1)M_{k-1}^s$ ($\exp(\gamma_3^{(a)})$)	3.51	[1.52,9.66]
$I(k > 1)(1 - M_{k-1}^s)S_{k-1}^{obs}$ ($\exp(\gamma_4^{(a)})$)	1.28	[0.72,2.28]
Moxifloxacin ($\exp(\gamma_5^{(a)})$)	1.07	[0.67,1.76]

[†] ∞ here means a big number

ratio as a function of α_0 and α_1 . The region is white indicates combinations of α_0 and α_1 where the lower bound of the 95% confidence interval is less than one. The green region indicates combinations of α_0 and α_1 where the null of no treatment difference is rejected in favor of moxifloxacin.

Inference would change relative to the benchmark assumption (circle in Figure 3), if say $\alpha_0 = 5.0, \alpha_1 = -3.0$ (triangle in Figure 3) or $\alpha_0 = -4.0, \alpha_1 = -10.0$ (square in Figure 3). At these combination of treatment-specific sensitivity analysis parameters, the estimated treatment effects are 2.19 (95% CI: 1.00-14.51) and 2.07 (95% CI: 0.97-5.38). To understand whether these combinations are “far” from the benchmark assumption, consider Figure 4. In the first row, we plot for each treatment group, the estimated distribution of time of culture conversion for these sensitivity analysis parameters (dashed and dotted lines) and the estimated distributions under the benchmark assumption (solid lines). In the second row, we plot for each

FIG 3. Contour plot of the estimated ratio of the odds of first becoming a culture converter at visit k given culture conversion at or after visit k , comparing moxifloxacin vs. ethambutol as a function of the sensitivity-analysis parameters α_0 and α_1 . The region is white indicates combinations of α_0 and α_1 where the lower bound of the 95% confidence interval is less than one. The green region indicates combinations of α_0 and α_1 where the null of no treatment difference is rejected in favor of moxifloxacin. Circle denotes the benchmark assumption ($\alpha_0 = 0.0, \alpha_1 = 0.0$). Triangle ($\alpha_0 = 5, \alpha_1 = -3$) and square ($\alpha_0 = -4, \alpha_1 = -10$) denotes two combinations discussed in the text.




TABLE 3
(Exponentiated) parameter estimates and 95% confidence intervals from the model for culture results. See $b_z(k, O_{k-1}, X; \gamma^{(b)})$ for form of the model.

Intercept	Odds	95% CI
Wk 1 ($\exp(\gamma_{0,0,1}^{(b)})$)	0.04	[0.02,0.08]
Wk 2 ($\exp(\gamma_{0,0,1}^{(b)})$)	0.03	[0.02,0.06]
Wk 3 ($\exp(\gamma_{0,0,1}^{(b)})$)	0.08	[0.04,0.14]
Wk 4 ($\exp(\gamma_{0,0,1}^{(b)})$)	0.10	[0.05,0.18]
Wk 5 ($\exp(\gamma_{0,0,1}^{(b)})$)	0.10	[0.05,0.17]
Wk 6 ($\exp(\gamma_{0,0,1}^{(b)})$)	0.24	[0.13-0.41]
Wk 7 ($\exp(\gamma_{0,0,1}^{(b)})$)	0.22	[0.12,0.40]
Wk 8 ($\exp(\gamma_{0,0,1}^{(b)})$)	0.49	[0.26,0.95]
Predictor	Odds Ratio	95% CI
$I(k > 1)M_{k-1}^c$ ($\exp(\gamma_1^{(b)})$)	4.00	[2.00,9.45]
$(k > 1)(1 - M_{k-1}^c)C_{k-1}^{obs}$ ($\exp(\gamma_2^{(b)})$)	6.37	[4.08,10.39]
$I(k > 1)M_{k-1}^s$ ($\exp(\gamma_3^{(b)})$)	5.16	[1.67,22.47]
$I(k > 1)(1 - M_{k-1}^s)S_{k-1}^{obs}$ ($\exp(\gamma_4^{(b)})$)	3.93	[2.51,6.18]
Moxifloxacin ($\exp(\gamma_5^{(b)})$)	2.06	[1.46,3.19]
Cavitation ($\exp(\gamma_6^{(b)})$)	1.16	[0.81,1.75]

treatment group, the signed Kolmogorov distance between the estimated distribution of time of culture conversion for given α and the estimated distribution function of time of culture conversion under the benchmark assumption. The signed Kolmogorov distance for treatment group z with sensitivity analysis parameter α_z equals $\widehat{F}_z(k_z^*; \alpha_z) - \widehat{F}_z(k_z^*; 0)$, where

$$k_z^* = \operatorname{argmax}_k \left| \widehat{F}_z(k; \alpha_z) - \widehat{F}_z(k; 0) \right|$$

and $\widehat{F}_z(k; \alpha_z)$ is the estimated cumulative distribution function. When $\alpha_0 = 5.0$ and $\alpha_1 = -3.0$, the signed distances for the ethambutol and moxifloxacin arms are 0.047 and -0.11 ; the latter being a fairly sizable difference (for the moxifloxacin arm, the estimated probability of culture conversion by Visit 5 is 49.6% under the benchmark assumption and 38.2% when $\alpha_1 = -3.0$). Further, the distances are of opposite signs (i.e., the bias is differential between arms). When we look at other combinations of sensitivity analysis parameters where the null hypothesis is not rejected, the sensitivity analysis parameter for the moxifloxacin arm is less than or equal to -3.0 and the associated signed distances are at least as extreme as -0.11 . When $\alpha_0 = -4.0$ and $\alpha_1 = -10.0$, the signed distances are -0.11 and -0.16 for the ethambutol and moxifloxacin arms, respectively. Here the signs are in the same direction, but the choice of sensitivity analysis parameters yield results that are very close to the worst case bounds that assume that all missing

TABLE 4

(Exponentiated) parameter estimates and 95% confidence intervals from the model for smear results. See $d_z(k, M_k^c, C_k^{obs}, O_{k-1}, X; \gamma^{(d)})$ for form of the model.


Intercept	Odds	95% CI
Wk 1 ($\exp(\gamma_{0,0,1}^{(d)})$)	0.25	[0.13, 0.42]
Wk 2 ($\exp(\gamma_{0,0,1}^{(d)})$)	0.34	[0.20,0.56]
Wk 3 ($\exp(\gamma_{0,0,1}^{(d)})$)	0.35	[0.21,0.58]
Wk 4 ($\exp(\gamma_{0,0,1}^{(d)})$)	0.21	[0.11,0.37]
Wk 5 ($\exp(\gamma_{0,0,1}^{(d)})$)	0.35	[0.19,0.65]
Wk 6 ($\exp(\gamma_{0,0,1}^{(d)})$)	0.20	[0.10,0.39]
Wk 7 ($\exp(\gamma_{0,0,1}^{(d)})$)	0.23	[0.12,0.43]
Wk 8 ($\exp(\gamma_{0,0,1}^{(d)})$)	0.24	[0.11,0.53]
Predictor	Odds Ratio	95% CI
M_k^c ($\exp(\gamma_1^{(d)})$)	1.24	[0.61, 2.61]
$(1 - M_k^c)C_k^{obs}$ ($\exp(\gamma_2^{(d)})$)	10.73	[5.58, 31.18]
$(1 - M_k^c)C_k^{obs} * Cav$ ($\exp(\gamma_3^{(d)})$)	0.46	[0.16,1.04]
$I(k > 1)M_{k-1}^c$ ($\exp(\gamma_4^{(d)})$)	1.52	[0.66,3.46]
$I(k > 1)(1 - M_{k-1}^c)C_{k-1}^{obs}$ ($\exp(\gamma_5^{(d)})$)	1.44	[0.92,2.38]
$I(k > 1)M_{k-1}^s$	3.66 ($\exp(\gamma_6^{(d)})$)	[1.45,14.53]
$I(k > 1)(1 - M_{k-1}^s)S_{k-1}^{obs}$ ($\exp(\gamma_7^{(d)})$)	6.99	[4.50,11.81]
Moxifloxacin ($\exp(\gamma_8^{(d)})$)	0.97	[0.63,1.48]
Cavitation ($\exp(\gamma_9^{(d)})$)	1.18	[0.75,1.85]

cultures are positive. From a clinical perspective, inferences relative to the benchmark assumption are fairly robust.

5. Discussion. Conde et al. (2009) did not compare the treatments with respect to time of culture conversion. Rather, they compared the treatment-specific probabilities of being a culture converter at or prior to week 8, which is equivalent to having a negative culture at week 8. They used two different methods. The primary method assumed that all missing cultures at week 8 were positive (moxifloxacin: 77.0%; ethambutol: 62.5%; difference: 14.5 percentage points, 95% CI [−0.0 percentage points, 29.0 percentage points]); the secondary method excluded patients who were missing their week 8 culture, assuming that the missing cultures were missing completely at random (moxifloxacin: 90.5%; ethambutol: 73.8%; difference: 16.7 percentage points, 95% CI [3.4 percentage points, 30.4 percentage points]). The former analysis is not statistically significant, while the latter analysis does suggest a statistically significant treatment effect in favor of moxifloxacin. In their analysis, no attempt was made to account for imbalances in baseline cavitation status between treatment groups.

It is tempting to think that this problem can be addressed by simply ana-

FIG 4. *First Row: Treatment-specific estimated distribution of time of culture conversion for the benchmark ($\alpha_0 = 0, \alpha_1 = 0$) and alternative ($\alpha_0 = 5.0, \alpha_1 = -3$) sensitivity-analysis parameters. Second Row: For each treatment group, the signed Kolmogorov distance between the estimated distribution of time of culture conversion for given α and the estimated distribution function of time of culture conversion under the benchmark assumption.*



lyzing the culture results using standard statistical methods for longitudinal binary data, e.g., marginal models and generalized linear mixed models. A marginal model, fit using generalized estimating equations, identifies, under the assumption that the culture results are missing completely at random, the probability of a negative culture at each visit k ; it does not admit identification of the distribution of time of culture conversion. In contrast, a generalized linear mixed model is a fully parametric model for the joint distribution of the culture results and, under the missing at random assumption, admits identification of the distribution of time of culture conversion. However, the modeling assumptions are too strong as they essentially allow the "imputation" of missing culture results that are not needed to identify the distribution of interest, e.g., the imputation of missing cultures that are preceded by positive cultures. Further, the model induces testable restrictions and, as discussed by [Robins and Gill \(1997\)](#), the missing at random assumption is often unrealistic in follow-up studies with intermittent missing data. Nonetheless, we fit a logistic-normal generalized linear mixed model to the culture data with fixed effects for time, treatment and cavitation, and a random intercept. Using the model, we estimated, within levels of time, treatment and cavitation, the induced probability of a negative culture among those with missing cultures. Many of the estimated probabilities were either greater than one or less than zero, suggesting inadequate model fit.

An alternative way of analyzing the culture data would treat the data for each patient as the set of times of culture conversion that are consistent with their observed culture data (i.e., the coarsening set) and estimate the distribution of time to culture conversion under the coarsening at random (CAR) assumption [[Gill, Van Der Laan and Robins \(1997\)](#); [Heitjan and Rubin \(1991\)](#); [Heitjan \(1993, 1994\)](#)]. This assumption states that the coarsening process provides no information about the time of culture conversion beyond conveying that the true event time is in the observed coarsening set. Under CAR, the coarsening process is "ignorable," i.e., it factors out of the likelihood for the observed data. Under this assumption, the estimated value of β is 2.92 (95%: 1.09-11.95). The result is statistically significant and favors moxifloxacin.

We have assumed, as in most analyses of culture conversion data, that the test results are measured without error. We know that this is not the case. It would be interesting to use known information about the sensitivity and specificity of the culture and smear procedures to learn about the "true" distribution of time of culture conversion. This will be the subject of future research.

In our analysis, we did not have access to the reasons for missingness. We know them to be a combination of three main sources: culture contamination, inability to produce sputum and skipped clinic visits. In the former two cases, the culture results are more likely to be negative. Contamination of sputum cultures with bacteria from the mouth and airways occurs in 2-10% of specimens and varies by laboratory. Patients producing smaller amounts of sputum that is mixed with saliva are more likely to have contamination; therefore, patients who have responded to therapy (i.e., have negative cultures) and no longer are producing large volumes of sputum may be more likely to have a contaminated specimen. Patients with treated tuberculosis who can no longer produce sputum are also likely to have responded to therapy and have negative cultures. If most of the missing data is due to these two causes, it is not so surprising that the results of the benchmark analysis are so close to the “best-case” bounds.

In summary, we introduced a novel benchmark assumption that allows us to “learn” about the distribution of just those unknown culture results that are absolutely necessary to identify the distribution of time of culture conversion by “borrowing strength” from patients as similar as possible (with respect to baseline cavitation status, treatment assignment, and observed culture and sputum results) and on whom the distribution of these culture results is identified. We evaluated the sensitivity of inferences to our benchmark assumption by embedding it within a class of model assumptions indexed by sensitivity-analysis parameters. While the sensitivity-analysis parameters themselves are not scientifically interpretable, the induced distribution of time of culture conversion (and functionals thereof) can be estimated and compared with that under the benchmark assumption. If the differences are judged “large” by scientific experts, we hope that they will comment on the fragility or robustness of the benchmark inference. Except in rare settings where the treatment effects are so dramatic or missing data are so minor, we see no alternatives to sensitivity analysis aided by scientific judgement.

Acknowledgements. The authors would like to thank Jonghyeon Kim, Chad Heilig, Malathi Ram, Pei-Jean Feng and Swarnadip Ghosh for assistance during the conduct of this research.

APPENDIX

A straightforward application of the law of total probability entails that under models (3.6) - (3.9),

$$P[T = j | \mathbf{O} = \mathbf{o}^{(j-1)}; \gamma] = \frac{g_{j-1}(0, \bar{O}_j, X; \gamma)}{\sum_{y=0}^1 g_{j-1}(y, \bar{O}_j, X; \gamma)}$$

where

$$\begin{aligned} g_j(y, \bar{O}_{j+1}, X; \gamma) = & (1 - \text{expit}\{a(j+1, O_j(0, y), X; \gamma^{(a)})\}) \times \\ & \text{expit}\{b(j, O_{j-1}, X; \gamma^{(b)})\}^y \times \\ & (1 - \text{expit}\{b(j, O_{j-1}, X; \gamma^{(b)})\})^{(1-y)} \times \\ & \text{expit}\{b(j+1, O_j(0, y), X; \gamma^{(b)})\} \times \\ & (1 - \text{expit}\{c(j, 0, y, O_{j-1}, X; \gamma^{(c)})\})^{(1-M_j^s)} \times \\ & \text{expit}\{c(j, 0, y, O_{j-1}, X; \gamma^{(c)})\}^{M_j^s} \times \\ & (1 - \text{expit}\{c(j+1, 0, 1, O_j(0, y), X; \gamma^{(c)})\})^{(1-M_{j+1}^s)} \times \\ & \text{expit}\{c(j+1, 0, 1, O_j(0, y), X; \gamma^{(c)})\}^{M_{j+1}^s} \times \\ & (1 - \text{expit}\{d(j, 0, y, O_{j-1}, X; \gamma^{(d)})\})^{(1-S_j)} \times \\ & \text{expit}\{d(j, 0, y, O_{j-1}, X; \gamma^{(d)})\}^{S_j} \times \\ & (1 - \text{expit}\{d(j+1, 0, 1, O_j(0, y), X; \gamma^{(d)})\})^{(1-S_{j+1})} \times \\ & \text{expit}\{d(j+1, 0, 1, O_j(0, y), X; \gamma^{(d)})\}^{S_{j+1}} \end{aligned}$$

and $O_j(0, y)$ represents observed data visit j with M_j^c set to 0 and C_j set y .

Notice that if $c(k, M_k^c, C_k^{obs}, O_{k-1}, X; \gamma^{(c)})$ does not depend on C_k^{obs} and C_{k-1}^{obs} for all k , then $P[T = j | \mathbf{O} = \mathbf{o}^{(j-1)}; \gamma]$ does not depend on $\gamma^{(c)}$.

REFERENCES

- AMERICAN THORACIC SOCIETY, (2000). Diagnostic standards and classification of tuberculosis in adults and children. *American Journal of Respiratory and Critical Care Medicine* **161** 1376-1395.
- CONDE, M. B., EFRON, A., LOREDO, C., DE SOUZA, G. R. M., GRAÇA, N. P., CEZAR, M. C., RAM, M., CHAUDHARY, M. A., BISHAI, W. R., KRITSKI, A. L. and CHAISSON, R. E. (2009). Moxifloxacin versus ethambutol in the initial treatment of tuberculosis: a double-blind, randomised, controlled phase II trial. *Lancet* **373** 1183-1189.
- COX, D. R. (1972). Regression models and life-tables (with discussion). *Journal of the Royal Statistical Society Series B* **34** 187-220.
- COX, D. and BARNDORFF-NIELSEN, O. (1994). *Inference and asymptotics*. Chapman & Hall/CRC.
- EUROPEAN MEDICINES AGENCY, COMMITTEE FOR MEDICINAL PRODUCTS FOR HUMAN USE, (2010). *Addendum to the Note for Guidance on Evaluation of Medicinal Products Indicated for Treatment of Bacterial Infections to Specifically Address the Clinical Development of New Agents to Treat Disease due to Mycobacterium Tuberculosis*. European Medicines Agency, London.
- GILL, R. D., VAN DER LAAN, M. J. and ROBINS, J. M. (1997). Coarsening at random: characterizations, conjectures and counter-examples. In *Proceedings of the First Seattle Symposium in Biostatistics: Survival Analysis* (D. Y. LIN and T. R. FLEMING, eds.) 255-294. Springer, Berlin.
- HEITJAN, D. F. (1993). Ignorability and coarse data: Some biomedical examples. *Biometrics* **49** 1099-1109.
- HEITJAN, D. F. (1994). Ignorability in general incomplete-data models. *Biometrika* **81** 701-708.
- HEITJAN, D. F. and RUBIN, D. B. (1991). Ignorability and coarse data. *Annals of Statistics* **19** 2244-2253.
- NEWBY, W. K. and MCFADDEN, D. (1994). Large sample estimation and hypothesis testing. In *Handbook of Econometrics*, (R. F. Engle and D. L. McFadden, eds.) **IV** 2111-2245. Elsevier, Amsterdam.
- ROBINS, J. M. and GILL, R. D. (1997). Non-response models for the analysis of non-monotone ignorable missing data. *Statistics in Medicine* **16** 39-56.

DANIEL SCHARFSTEIN
 ADIAN McDERMOTT
 DEPARTMENT OF BIostatISTICS
 JOHNS HOPKINS BLOOMBERG SCHOOL OF PUBLIC HEALTH
 615 NORTH WOLFE STREET
 BALTIMORE, MARYLAND 21205
 E-MAIL: dscharf@jhsph.edu
amcdermo@jhsph.edu

ANDREA ROTNITZKY
 DEPARTMENT OF ECONOMICS
 UNIVERSIDAD TORCUATO DI TELLA
 SAENZ VALIENTE 1010
 1428 BUENOS AIRES
 E-MAIL: arotnitzky@utdt.edu

MARIA ABRAHAM
 STATISTICS COLLABORATIVE
 1625 MASSACHUSETTS AVE NW
 SUITE 600
 WASHINGTON, DC 20036
 E-MAIL: maria@statcollab.com

RICHARD CHAISSON
 JOHNS HOPKINS CENTER FOR TUBERCULOSIS RESEARCH
 1550 ORLEANS ST., 1M.08
 BALTIMORE, MD 21231
 E-MAIL: rchaiss@jhmi.edu

LAWRENCE J. GEITER
OTSUKA NOVEL PRODUCTS-TB
OTSUKA PHARMACEUTICAL DEVELOPMENT AND COMMERCIALIZATION, INC.
2440 RESEARCH BOULEVARD
ROCKVILLE, MD 20850
E-MAIL: Lawrence.Geiter@otsuka-us.com